

Home Isolation Instructions for People with Coronavirus Disease (COVID-19) Infection

The following instructions are for people who have COVID-19. It also includes information for their families or caregivers.


Home Care

There is no specific treatment for the virus that causes COVID-19. Here are steps that you can take to help you get better:

- Rest
- Drink plenty of fluids
- Take acetaminophen (Tylenol®) to reduce fever and pain.
- Note that children younger than age 2 should not be given any over-the-counter cold medications without first speaking with a doctor.
- Note that these medicines do not “cure” the illness and do not stop you from spreading germs.

Seeking Medical Care

Seek prompt medical care if your symptoms get worse, especially if you are at a higher risk of serious illness. This includes people who are age 65 years and older, pregnant, or have a health problem such as a chronic disease or a weak immune system. It is recommended that you seek medical care for serious symptoms, such as:



People with life-threatening symptoms should call 911. Tell the dispatch personnel that you have COVID-19. If possible, put on a facemask before emergency medical services arrive.

If it's not urgent, call ahead before visiting your doctor, you may be able to get advice by phone. If you do visit a healthcare facility, put on a mask before you enter to protect others from catching the infection from you.

COVID-19 may be stressful for people, visit <http://www.stancounty.com/bhrs/covid.shtm> to learn how to care for your mental health and support your loved ones. If you need to speak with someone about your mental health, contact your doctor or call (209) 558-4600 for Stanislaus County Emergency Support, available 24/7. If you speak another language, language assistance services, free of charge, are available to you. Call 1-888-376-6246 (TTY: 711).

For more information on COVID-19:

Call (209) 558 7535 OR Visit <http://www.schsa.org/coronavirus/>

Adapted from Los Angeles County Department of Public Health - Revised 12.03.20

Home Isolation Instructions for People with Coronavirus Disease (COVID-19) Infection

PROTECTING OTHERS

Follow the steps below to help prevent the disease from spreading to people in your home and your community.

Stay home except to get medical care

- Do not go to work, school, or public areas.
- If you must leave home while you are sick, do not use public transportation. Use a personal vehicle if possible. If you cannot drive yourself, keep as much distance as possible between you and the driver, leave the windows down and wear a mask.
- If you do not have someone to help you, if possible, arrange for food and other necessities to be left at your door. If you need to meet someone at your door, wear a mask.
- Stay home until at least 10 days have passed after your symptoms first appeared AND at least **1 day** after you have recovered. Recovery means that your fever is gone for **24 hours** without the use of fever-reducing medications and your symptoms have improved.

Separate yourself from other people in your home

- Stay in a specific room and away from other people in your home as much as possible. It is particularly important to stay away from people who are at higher risk of serious illness. Consider alternate living arrangements for them if at all possible.
- Use a separate bathroom. If this is not possible, clean the bathroom after use (see below)
- Stay at least 6 feet from others.
- Open windows or use a fan or an air conditioner in shared spaces in the home, if possible, to ensure good airflow.
- Do not allow visitors and limit the number of people in your home.
- Do not handle pets or other animals while you are sick.
- Do not prepare or serve food to others.
- Do not care for children if possible.

Wear a facemask when you are around others

You should wear a facemask when you are around other people (e.g., sharing a room or vehicle) or pets and before you enter a hospital or doctor's office. If you are not able to wear a facemask (for example, because it causes trouble breathing), then people who live with you should not be in the same room with you. If they must enter your room, they should wear a facemask. After leaving your room, they should immediately wash their hands, remove and dispose of their facemask, and then wash their hands again.

Cover your coughs and sneezes

Cover your mouth and nose with a tissue when you cough or sneeze. Throw used tissues in a lined trash can; immediately wash your hands.

Avoid sharing personal household items

Do not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people or pets in your home. Wash them thoroughly with soap and water after use.

For more information on COVID-19:

Call (209) 558 7535 OR Visit <http://www.schsa.org/coronavirus/>

Adapted from Los Angeles County Department of Public Health - Revised 12.03.20

Home Isolation Instructions for People with Coronavirus Disease (COVID-19) Infection

Clean your hands often

Wash your hands often and thoroughly, especially after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food. Use soap and water for at least 20 seconds. If soap and water are not available, clean your hands with a hand sanitizer that contains at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Use soap and water if your hands are visibly dirty.

Clean and disinfect all “high-touch” surfaces every day

High touch surfaces include counters, tabletops, doorknobs, bathroom fixtures, toilets, phones, keyboards, tablets, and bedside tables. Also, clean and disinfect any surfaces that may have body fluids on them. Use household cleaning and disinfectant sprays or wipes, according to the product label instructions. See cleaning instructions in *Cleaning And Disinfecting Your Home* on the [Public Health website](#).

How long do I need to isolate from others?

Stay home and away from others until at least 10 days have passed after your symptoms first appeared AND at least **1 day** after you have recovered. Recovery means that your fever is gone for **24** hours without the use of fever-reducing medications and your symptoms have improved.

INFORMATION FOR YOUR CLOSE CONTACTS

Quarantine

Be sure to tell all of your close contacts that they need to be in quarantine for **10 days** after their last contact with you. Close contacts include all household members, any intimate contacts, and all individuals who were within 6 feet of you for more than 15 minutes, starting 48 hours before your symptoms began until your isolation period ends. In addition, anyone who had contact with your body fluids and/or secretions (such as were coughed on/sneezed on, shared utensils or saliva or provided care to you without wearing protective equipment) needs to be in quarantine. They should self-quarantine even if they feel well because it can take 2– 14 days for them to show symptoms. See the [Home quarantine guidance for those exposed to COVID-19 \(English\)](#), [\(Spanish\)](#).

Precautions for close contacts

It is recommended that everyone stays at least 6 feet away from you while you are under home isolation. If this is not possible, anyone who continues to be in close contact with you will need to extend their quarantine period to **10 days** from the last time they had close contact with you.

Your caregivers and household contacts should wear a disposable facemask and gloves if they clean your room or bathroom or come into contact with your body fluids, and/or secretions (such as sweat, saliva, sputum, nasal mucus, vomit, urine, or diarrhea). They should remove and dispose of their gloves first, clean their hands, then remove and dispose of their facemask, and clean their hands again.

What should I do if I have additional questions?

Visit the Public Health website [for more information](#) about coronavirus. Please call your health care provider for any questions related to your health.

For more information on COVID-19:

Call (209) 558 7535 OR Visit <http://www.schsa.org/coronavirus/>

Adapted from Los Angeles County Department of Public Health - Revised 12.03.20